REFERENCES/BIBLIOGRAPHY

1] Aoki, J., & Uchino, K. (2011). Treatment of Risk Factors to Prevent Stroke, 463–474. https://doi.org/10.1007/s13311-011-0054-0

2] Amen, M. R. (2016). Assessment of hypertensive patients' knowledge about lifestyle risk factors and warning signs of stroke. Journal of Contemporary Medical Sciences, 2(5), 28-32.

3] Banerjee, T. K., Roy, M. K., &Bhoi, K. K. (2005). Is stroke increasing in India--preventive measures that need to be implemented. Journal of the Indian Medical Association, 103(3), 162-164.

4] Becker, K. J., Fruin, S., Gooding, D., Tirschwell, D. L., Love, J., & Mankowski, T. M. (2001). Community-Based Education Improves Stroke Knowledge, 2499, 34–43.

5] Castilla-guerra, L., & Fernández-moreno, C. (2012). Update on the Management of Hypertension for Secondary Stroke, 1–7. https://doi.org/10.1159/000336836.

6] Das, P., Manimay, B., Paul, R., & Singh, R. (2010). Study of underlying risk factors for stroke in young. Biomedical research, 21(3).

7] Ellis, C. (2013). Poststroke Knowledge and Symptom Awareness : A Global Issue for Secondary stroke prevention. 572–581. https://doi.org/10.1159/000351209

8] Franklin, B. A., & Sanders, W. (2000). Reducing the risk of heart disease and stroke. The Physician and sportsmedicine, 28(10), 19-26.

9] Gourie-Devi, M. (1987). Can India afford neuroepidemiology?. NeurolIndia, 35, 125-127.

10] George, M. C., Tong, X., & Bowman, B. A. Prevalence of Cardiovascular Risk Factors and Strokes in Younger Adults. JAMA Neurol. 2017; 74: 695–703.

11] Green, B. T., Haley, E., Eliasziw, M., & Hoyte, K. (2007). Education in stroke prevention : Efficacy of an educational counselling intervention to increase knowledge in stroke survivors, 29(2).

12] Gutie, E., Escamilla-garza, J. M., Villarreal, J., Investigators, G., Carranco-chavez, D., Chapa-rodríguez, R., & Roy, M. (2011). Knowledge of Ischemic Stroke Risk Factors and Warning Medical Students. https://doi.org/10.1161/STROKEAHA.110.597062

13] Grau, A. J., Barth, C., Geletneky, B., Ling, P., Palm, F., Lichy, C. &Buggle, F.

(2009). Association between recent sports activity, sports activity in young adulthood, and stroke. Stroke, 40(2), 426-431.
14]Haapaniemi, H., Hillborn M, Juvela S (1997) Lifestyle associated risk factors for acute brain infraction in young adults among persons of working age. Stroke. Jan,28(1):26-30
15]Jain, S., &Maheshwari, M. C. (1986). Cerebrovascular diseases: a review of the Indian experience in the last 35 years. Neuroepidemiology, 5(1), 1-16.
16] Jørgensen, H. S., Nakayama, H., Raaschou, H. O., & Olsen, T. S. (1995). Intracerebral hemorrhage versus infarction: stroke severity, risk factors, and prognosis. Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society, 38(1), 45-50.
17]Kotrabasappa, K., &Deepadarshan, H. (2016). Risk factors and outcome of stroke in young in a tertiary care hospital.
18]Nakibuuka Jane MBChB (MAK), Prof . Elly Katabira , Prof . Anthony Furlan ,et. al. Proposal submitted in fulfilment of the requirement for the award of Doctor of Philosophy of Makerere University Kampala , Uganda Version 2. (2011).Improving stroke prevention and outcomes in uganda : populationsurvey and hospital based study.
19] Müller-Nordhorn, J., Nolte, C. H., Rossnagel, K., Jungehülsing, G. J., Reich, A., Roll, S. &Willich, S. N. (2006). Knowledge about risk factors for stroke: A population-based survey with 28 090 participants. Stroke, 37(4), 946-950.
20]Kaul S.,Bandarn V C et al (2009) Stroke burden and risk factors in developing countries with special reference to India. J Indian Med Assoc. June;107(6):358,367-370
21]Kumar, D., Singh, R. P., Hooda, A. K., Deswal, B. S., & Hassan, K. M. (2018). Lifestyle riskfactors of stroke in young Indians, *5*(March), 2–9. https://doi.org/10.18231/2394-2738.2018.0002
22]Kristensen, B., Malm, J., Carlberg, B., Stegmayr, B., Backman, C., Fagerlund, M., & Olsson, T. (1997). Epidemiology and etiology of ischemic stroke in young adults aged 18 to 44 years in northern Sweden. Stroke, 28(9), 1702-1709.
23]Lakka, T. A., &Laaksonen, D. E. (2007). Physical activity in prevention and treatment of the metabolic syndrome. Applied physiology, nutrition, and metabolism, 32(1), 76-88.
24] Law, M. R., Wald, N. J., Wu, T., Hackshaw, A., & Bailey, A. (1994). Systematic underestimation of association between serum cholesterol concentration and ischaemic heart disease in observational studies: data from the BUPA study. Bmj, 308(6925), 363-

366.
25]Lee, C. D., Folsom, A. R., & Blair, S. N. (2003). Physical activity and stroke risk: a meta-analysis. Stroke, 34(10), 2475-2481.
26]Lee, I. M., Hennekens, C. H., Berger, K., Buring, J. E., & Manson, J. E. (1999). Exercise and risk of stroke in male physicians. Stroke, 30(1), 1-6.
27]Lipska, K., Sylaja, P. N., Sarma, S. P., Kutty, V. R., Thankappan, K. R., Vasan, R. S., &Radhakrishnan, K. (2007). Risk factors for acute ischaemic stroke in young adults in South India. Journal of Neurology, Neurosurgery & Psychiatry.
28] Lopez, G. A., Afshinnik, A., & Samuels, O. (2011). Care of the stroke patient: routine management to lifesaving treatment options. *Neurotherapeutics*, *8*(3), 414.
29]Marwat, M. A., Usman, M., & Hussain, M. (2009). Stroke and its relationship to risk factors.Gomal j Med Sci,7(1):17-19
30]Mishra, N. K., &Khadilkar, S. V. (2010). Stroke program for India. Annals of Indian Academy of Neurology, 13(1), 28
31]Nakibuuka, J., Sajatovic, M., Katabira, E., Ddumba, E., Byakika-tusiime, J., & Furlan, A. J. (2014). Knowledge and Perception of Stroke : A Population-Based Survey in Uganda, *2014*.
32] Nedeltchev, K., der Maur, T. A., Georgiadis, D., Arnold, M., Caso, V., Mattle, H. P., ...& Baumgartner, R. W. (2005). Ischaemic stroke in young adults: predictors of outcome and recurrence. Journal of Neurology, Neurosurgery & Psychiatry, 76(2), 191-195.
33]Nayak, S. D., Nair, M., Radhakrishnan, K., &Sarma, P. S. (1997). Ischaemic stroke in the young adult: clinical features, risk factors and outcome. The National medical journal of India, 10(3), 107-112.
34]Nicol, M. B., & Thrift, A. G. (2005). Knowledge of risk factors and warning signs of stroke. *Vascular health and risk management*, *1*(2), 137.
35]Prasad, K., &Singhal, K. K. (2010). Stroke in young: an Indian perspective. *Neurology India*, *58*(3), 343
36]Pandian, J. D., Jaison, A., Deepak, S. S., Kalra, G., Shamsher,S., Lincoln, D. J., & Abraham, G. (2005). Public awareness of warning symptoms, risk factors, and treatment of strokein northwest India. *Stroke*, *36*(3), 644-648.
37]Rexrode, K. M., Hennekens, C. H., Willett, W. C., Colditz, G. A., Stampfer, M. J., Rich-Edwards, J. W., ...& Manson, J. E. (1997). A prospective study of body mass

index, weight change,and risk of stroke in women. *Jama*, *277*(19), 1539-1545.
38] Rohr, J., Kittner, S., Feeser, B., Hebel, J. R., Whyte, M. G., Weinstein, A. &Macko, R. (1996). Traditional risk factors and ischemic stroke in young adults: the Baltimore-Washington Cooperative Young Stroke Study. *Archives of neurology*, *53*(7), 603-607.
39]Sacco, R. L., Gan, R., Boden-Albala, B., Lin, I. F., Kargman, D. E., Hauser, W. A., ...& Paik, M. C. (1998). Leisure-time physical activity and ischemic stroke risk: the Northern Manhattan Stroke Study. *Stroke*, *29*(2), 380-387
40] Sacco, R. L., Adams, R., Albers, G., Alberts, M. J., Benavente, O., Furie, K., ...& Johnston, S. C. (2006). Guidelines for prevention of stroke in patients with ischemic stroke or transient ischemic attack: a statement for healthcare professionals from the American Heart Association/American Stroke Association Council on Stroke: co-sponsored by the Council on Cardiovascular Radiology and Intervention: the American Academy of Neurology affirms the value of this guideline. *Circulation*, *113*(10), e409-e449.
41] Saengsuwan, J., Suangpho, P., & Tiamkao, S. (2017). Knowledge of Stroke Risk Factors and Warning Signs in Patients with Recurrent Stroke or Recurrent Transient Ischaemic Attack inThailand, *2017*.
42] Schneider, S., Kornejeva, A., Vibo, R., & Kõrv, J. (2017). Risk Factors and Etiology of Young Ischemic Stroke Patients in Estonia, *2017*.
43] Shinton, R., &Beevers, G. (1989). Meta-analysis of relation between cigarette smoking and stroke. *Bmj*, *298*(6676), 789-794.
44] Singhal, A. B., Fullerton, H. J., Jauch, E. C., & Kittner, S. J. (2013). Recognition and management of stroke in young adults and adolescents.
45] Zulkifly MFM, Ghazali SE et al (2016), A review of risk factors for cognitive impairment in stroke survivors; The Scientific World Journal Volume 2016, Article ID 3456943, 16 pageshttp://dx.doi.org/10.1155/2016/3456943